

Math 365 – Wednesday 2/6/19
Section 2.4: Sequences and summations

Exercise 12.

- (a) For each of the following sequences, compute the terms a_0 , a_1 , a_2 , and a_3 .
- (a) $a_n = 3$;
 - (b) $a_n = 7 + 4^n$;
 - (c) $a_n = 2^n + (-2)^n$.
- (b) For each of the following sequences defined by recurrence relations and initial conditions, answer the following.
- (a) Compute the first four terms (a_0 , a_1 , a_2 , and a_3).
 - (b) Decide if $\{a_n\}$ is arithmetic, geometric, or neither. If it is arithmetic or geometric, then find a closed formula for a_n .
 - (i) The sequence satisfying $a_0 = 2$ and $a_n = \frac{1}{2}a_{n-1}$.
 - (ii) The sequence satisfying $a_0 = -1$ and $a_n = a_{n-1} + 5$.
 - (iii) The sequence satisfying $a_0 = 1$, $a_1 = -1$ and $a_n = a_{n-2} * a_{n-1}$.
 - (iv) The sequence satisfying $a_0 = 2$ and $a_n = -a_{n-1}$.
- (c) Find at least three different sequences beginning with the terms 3, 5, 7 whose terms are generated by a simple formula or rule.
- (d) For the following sequences, try to find the pattern. Decide if they are arithmetic, geometric, or neither. If it's arithmetic or geometric, find a closed formula expressing the n th term of the sequence.
- (i) 5, 1, -3, -7, -11, ...
 - (ii) 1, 4, 9, 16, 25, ...
 - (iii) 3, 9, 27, 81, 243, ...
- (e) Show that both of the following sequences are solutions to the recurrence relation $a_n = -3a_{n-1} + 4a_{n-2}$ with initial condition $a_0 = 1$.
- (i) $a_n = 1$;
 - (ii) $a_n = (-4)^n$.

Exercise 13. (a) Compute the first three partial sums for the following infinite series.

- (i) $\sum_{i=0}^{\infty} 5i + 1$;
 - (ii) $\sum_{i=4}^{\infty} i(i + 1)$;
- (b) Calculate the following.
- (i) $\sum_{j=0}^8 (1 + (-1)^j)$
 - (ii) $\sum_{j=-1}^2 \sum_{i=2}^3 (2i + 3j)$
 - (iii) $\sum_{j \in \{-1, 4, 15\}} 2$

$$(iv) \sum_{j \in \{z \in \mathbb{Z} \mid |z| \leq 2\}} j$$

$$(v) \sum_{n=2}^5 a_n - a_{n-1} \text{ where } a_n = n!$$

$$(vi) \sum_{i=1}^{2000} i$$

$$(vii) \sum_{i=0}^{10} \frac{2}{3^i}$$

$$(viii) \sum_{i=0}^{\infty} \frac{2}{3^i}$$

Exercise 14. A little more.

(a) For each of these sequences find a recurrence relation satisfied by this sequence. (The answers are not unique because there are infinitely many different recurrence relations satisfied by any sequence.)

(i) $a_n = 2n + 3;$

(ii) $a_n = 5^n;$

(iii) $a_n = n^2.$

(b) Use partial sums to explain why, for any sequence a_0, a_1, \dots, a_n , that

$$\sum_{i=1}^n a_i - a_{i-1} = a_n - a_0.$$

[Let $S_n = \sum_{i=1}^n a_i - a_{i-1}$. Write out S_1, S_2, S_3 , and so on until you see the pattern. Then use the fact that $S_n = S_{n-1} + (a_n - a_{n-1}).$]

(c) Read in section 2.4 about product notation $\prod_{i=m}^n a_i$. Then, what are the values of the following products?

(a) $\prod_{i=0}^{10} i$

(b) $\prod_{i=5}^8 i(i+1)$

(c) $\prod_{i=1}^{100} (-1)^i$

(d) Express $n!$ using product notation.

Sequences

A *sequence* is a function a from a subset of the set of integers (usually $\mathbb{Z}_{\geq 0}$ or $\mathbb{Z}_{> 0}$) to a set S ,

$$a : \mathbb{Z}_{\geq 0} \rightarrow S \quad \text{or} \quad a : \mathbb{Z}_{> 0} \rightarrow S.$$

We write $a_n = a(n)$, and call a_n the *n th term of the sequence*.

Example

The sequence defined by the function

$$a : \mathbb{Z}_{> 0} \rightarrow \mathbb{Q} \quad \text{defined by} \quad n \mapsto 1/n$$

is the sequence

$$1, 1/2, 1/3, 1/4, \dots$$

We write $a_n = 1/n$.

We can also write such a sequence like

$$\{a(n)\}_{n=1,2,\dots} \quad \text{or} \quad \{a(n)\}_{n \in \mathbb{Z}_{> 0}}.$$

For example, the sequence above is $\{1/n\}_{n \in \mathbb{Z}_{> 0}}$.

Some different kinds of sequences

A *geometric sequence* (or *progression*) is a sequence of the form

$$c, cr, cr^2, cr^3, \dots, \quad \text{i.e.} \quad a : \mathbb{Z}_{\geq 0} \rightarrow S \quad \text{by} \quad n \mapsto cr^n,$$

for some constants c and r . (This is a discrete version of the exponential function $f(x) = cr^x$.)

An *arithmetic progression* is a sequence of the form

$$b, b + m, b + 2m, b + 3m, \dots, \\ \text{i.e.} \quad a : \mathbb{Z}_{\geq 0} \rightarrow S \quad \text{by} \quad n \mapsto b + mn,$$

for some constants b and m . (This is a discrete version of the linear function $f(x) = b + mx$.)

Notice, with a geometric sequence, the *ratio is constant*:

$$\text{if } a_n = cr^n, \quad \text{then } a_n/a_{n-1} = r \text{ for all } n.$$

And with an arithmetic sequence the *difference is constant*:

$$\text{if } a_n = b + mn, \quad \text{then } a_n - a_{n-1} = m \text{ for all } n.$$

(This is how we test to see if a sequence is geometric or arithmetic!)

Recurrence relations

A *recurrence relation* for a sequence is an equation that expresses a_n in terms of one or more of the previous terms of the sequence. For example:

$$\begin{aligned}a_n &= a_{n-1} * 2; \\a_n &= a_{n-2} + 1; \\a_n &= a_{n-1} + a_{n-2}.\end{aligned}$$

A sequence is called a *solution* to a recurrence relation if its terms satisfy the recurrence relation. For example,

$a_n = 3 * 2^n$ is a solution to the recurrence relation $a_n = a_{n-1} * 2$;
 $a_n = -2^n$ is also a solution to the recurrence relation $a_n = a_{n-1} * 2$;
 $a_n = c * 2^n$ is also a solution to the recurrence relation $a_n = a_{n-1} * 2$,
for any $c \in \mathbb{R}$.

An *initial condition* is a specified value for some fixed a_i (usually a_0 and/or a_1). Without initial conditions, there are usually many solutions to a recurrence relation. For example,

$a_n = 3 * 2^n$ is the only solution to the r. rel. $a_n = a_{n-1} * 2, a_0 = 3$.

Recurrence relations

A *recurrence relation* for a sequence is an equation that expresses a_n in terms of one or more of the previous terms of the sequence.

A sequence is called a *solution* to a recurrence relation if its terms satisfy the recurrence relation. For example,

$a_n = 3 * 2^n$ is a solution to the recurrence relation $a_n = a_{n-1} * 2$;
 $a_n = -2^n$ is also a solution to the recurrence relation $a_n = a_{n-1} * 2$;
 $a_n = c * 2^n$ is also a solution to the recurrence relation $a_n = a_{n-1} * 2$,
for any $c \in \mathbb{R}$.

And *initial condition* is a specified value for some fixed a_i (usually a_0 and/or a_1). Without initial conditions, there are usually many solutions to a recurrence relation. For example,

$a_n = 3 * 2^n$ is the only solution to the r. rel. $a_n = a_{n-1} * 2, a_0 = 3$.

A *closed formula* for a recurrence relation is a formula generating the sequence. We call a closed formula that satisfies a recurrence relation a *solution* to that relation. (Ex: $a_n = c * 2^n$)

Going from a recurrence relation to a closed form is like calculating integrals—it is not always even possible, let alone deterministic. We learn to recognize familiar types, and look for patterns.

Geometric: If $a_n = ra_{n-1}$, then

$$a_1 = r \cdot a_0, \quad a_2 = ra_1 = r(ra_0) = r^2a_0, \\ a_3 = ra_2 = r(r^2a_0) = r^3a_0 \dots$$

Claim: In general, $a_n = a_0r^n$ for whatever constant a_0 is.

Arithmetic: $a_n = m + a_{n-1}$, then

$$a_1 = m + a_0, \quad a_2 = m + a_1 = m + (m + a_0) = 2m + a_0, \\ a_3 = m + a_2 = m + (2m + a_0) = 3m + a_0 \dots$$

Claim: In general, $a_n = nm + a_0$ for whatever constant a_0 is.

Going from a recurrence relation to a closed form is like calculating integrals—it is not always even possible, let alone deterministic. We learn to recognize familiar types, and look for patterns.

Factorial: For $n \in \mathbb{Z}_{>0}$, we define *n factorial*, denoted $n!$, by

$$n! = n(n-1)(n-2) \cdots 2 \cdot 1.$$

For example, $4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24$.

For convenience, we define $0! = 1$.

Then if $a_n = na_{n-1}$, we have

$$a_1 = 1 \cdot a_0, \quad a_2 = 2a_1 = 2(1 \cdot a_0) = (2 \cdot 1)a_0, \\ a_3 = 3a_2 = 3((2 \cdot 1)a_0) = (3 \cdot 2 \cdot 1)a_0 \dots$$

Claim: In general, $a_n = n!a_0$ for whatever constant a_0 is.

You try: Exercise 12

Summations

Recall, for a sequence $\{a_n\}$, the *summation notation*

$$\sum_{i=k}^{\ell} a_i = a_k + a_{k+1} + \cdots + a_{\ell}$$


and

$$\sum_{i=k}^{\infty} a_i = a_k + a_{k+1} + \cdots = \lim_{\ell \rightarrow \infty} \sum_{i=k}^{\ell} a_i.$$

For example, let $a_i = i$. Define $S_n = \sum_{i=1}^n a_i$. Then

$$S_1 = 1, \quad S_2 = 1 + 2 = 3, \quad S_3 = 1 + 2 + 3 = 6, \dots$$

Claim: In general, $S_n = \frac{n(n+1)}{2}$. "closed formula"




So $\sum_{i=1}^{\infty} a_i$ is not defined (the series does not converge).

On the other hand, let $a_i = (1/2)^i$, and define $S_n = \sum_{i=1}^n a_i$. Then

$$S_1 = 1/2, \quad S_2 = 1/2 + 1/4 = 3/4,$$

$$S_3 = 1/2 + 1/4 + 1/8 = 7/8 = 1 - 1/8,$$

$$S_4 = 1/2 + 1/4 + 1/8 + 1/16 = 15/16 = 1 - 1/16 \dots$$



Claim: In general, $S_n = 1 - \frac{1}{2^n}$. "closed formula"

So

$$\sum_{i=1}^{\infty} a_i = \lim_{n \rightarrow \infty} \left(1 - \frac{1}{2^n} \right) = 1.$$

Solving using partial sums

The finite sum

$$S_n = \sum_{i=0}^n a_i = a_0 + a_1 + \cdots + a_n$$

is called the *partial sum* for the series $S = \sum_{i=0}^{\infty} a_i$. We define

$S = \lim_{n \rightarrow \infty} S_n$. So to solve for S , it would be *very helpful* to get a closed form for S_n .

Example

Show

$$\sum_{i=0}^n cr^i = \begin{cases} c \left(\frac{r^{n+1}-1}{r-1} \right) & \text{if } r \neq 1 \\ c(n+1) & \text{if } r = 1 \end{cases}$$

using partial sums. Namely, show $rS_n = S_n + c(r^{n+1} - 1)$ and solve for S_n . Then calculate $\sum_{i=0}^{\infty} cr^i$.

Identities:

$$\sum_{i \in S} a_i + b_i = \sum_{i \in S} a_i + \sum_{i \in S} b_i \quad (\text{addition is commutative})$$

$$\sum_{i \in S} c * a_i = c * \sum_{i \in S} a_i \quad (\text{distributive property})$$

Set summations.

$$\sum_{a \in A} a \quad \text{means add up everything in } A.$$

$$\sum_{a \in A} f(a) \quad \text{means add up } f(a) \text{ for everything in } A.$$

Example:

$$\sum_{i \in \{2,4,6\}} i^2 = 2^2 + 4^2 + 6^2.$$

More notation

Double summations.

For example,

$$\begin{aligned}\sum_{i=1}^3 \sum_{j=i}^4 ij &= \sum_{i=1}^3 \left(\sum_{j=i}^4 ij \right) \\ &= \left(\sum_{j=1}^4 1 \cdot j \right) + \left(\sum_{j=2}^4 2 \cdot j \right) + \left(\sum_{j=3}^4 3 \cdot j \right) \\ &= (1 \cdot 1 + 1 \cdot 2 + 1 \cdot 3 + 1 \cdot 4) \\ &\quad + (2 \cdot 2 + 2 \cdot 3 + 2 \cdot 4) + (3 \cdot 3 + 3 \cdot 4).\end{aligned}$$

More special summations.

Theorem

We have the following special summation identities:

$$\begin{aligned}\sum_{i=1}^n i &= n(n+1)/2, \text{ and} \\ \sum_{i=0}^{\infty} ar^i &= \frac{a}{1-r} \quad \text{for } r \in (-1, 1).\end{aligned}$$

Notice

$$\sum_{i=1}^{\infty} i \cdot x^{i-1} = \frac{d}{dx} \sum_{i=0}^{\infty} x^i.$$

What can we conclude?

You try: Exercise 13